

# Ethnic Differences In The Pathogenesis of Obstructive Sleep Apnoea

Tahrani, Abd A.

DOI:

[10.1111/resp.13057](https://doi.org/10.1111/resp.13057)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Tahrani, AA 2017, 'Ethnic Differences In The Pathogenesis of Obstructive Sleep Apnoea: Exploring Non-Anatomical Factors', *Respirology*. <https://doi.org/10.1111/resp.13057>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

This is the peer reviewed version of the following article: [FULL CITE], which has been published in final form at [Link to final article using the DOI]. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Ethnic Differences In The Pathogenesis of Obstructive Sleep Apnoea: Exploring Non-Anatomical Factors

Abd A Tahrani

Institute of Metabolism and Systems Research, University of Birmingham, UK.

Centre of Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, UK

Department of Diabetes and Endocrinology, Birmingham Heartlands Hospital, Birmingham, UK

Corresponding author:

Abd A Tahrani

Department of Diabetes and Endocrinology,

Birmingham Heartlands Hospital,

Birmingham B9 5SS

07801549960

[a.a.tahrani@bham.ac.uk](mailto:a.a.tahrani@bham.ac.uk) or [abd.tahrani@nhs.net](mailto:abd.tahrani@nhs.net)

Key words: sleep apnoea, OSA, arousal threshold, Chinese, Caucasian, ethnicity

Obstructive sleep apnoea (OSA) is a common disorder that affects 17-26% of men and 9-28% of women<sup>1</sup>. The instability and increased collapsibility of the upper airways during sleep are the cardinal features of OSA, which result in recurrent upper airway obstruction resulting in either complete or partial cessation of airflow that leads to the characteristic features of OSA such as recurrent oxygen desaturations, disruption of sleep architecture and cyclical changes in heart rate, blood pressure and sympathetic activity<sup>2</sup>.

Considering that obesity is a major risk factor of OSA, it is not surprising that the prevalence of OSA increases with increasing prevalence of obesity<sup>3</sup>. OSA prevalence and severity have been shown to increase with weight gain and improve with weight loss<sup>4;5</sup>. In addition to obesity, craniofacial features and surface anatomy (such as cervicomentral angle and face width) and upper airway structures (such as tongue volume and soft palate length) are associated with OSA<sup>6-8</sup>. Furthermore, a crowded posterior oropharynx and a steep thyromental plane are associated with OSA in Asian and White patients independently of obesity<sup>9</sup>.

In addition to obesity and craniofacial anatomy, other non-anatomical factors such as low arousal thresholds (ArTH), passive critical closing pressure of the upper airway, loop gain, and upper airway muscle responsiveness play an important role in the pathogenesis of OSA<sup>10</sup>. The upper airway obstruction that occurs in OSA leads to an increase in CO<sub>2</sub> levels that leads to increased ventilatory drive and increasingly negative pharyngeal pressure which in turn can activate the upper airway dilator muscles to open the obstructed upper airway and restores its patency<sup>11</sup>. However, early arousals due to low ArTH can interrupt this process and result in restoration of upper airway patency before the development of adequate ventilatory drive, which can lead to further multiple upper airway obstructions<sup>11</sup>. A low respiratory ArTH is present in a third of patients with OSA<sup>11</sup>.

Studies that examine ethnic differences in OSA prevalence show inconsistent results<sup>12-14</sup>. However, several studies have shown that Asian patients have similar OSA prevalence to that reported in studies conducted in Europe and North America. This is despite the fact that Asians were less obese

(based on body mass index (BMI))<sup>15</sup>. In addition, the OSA phenotype differs amongst ethnicities, which can partly be explained by differences in obesity and craniofacial anatomy. Our group has shown that differences in adiposity, particularly upper body fat distribution, may explain our findings of a higher prevalence of OSA in White Europeans vs. South Asians with Type 2 diabetes<sup>16</sup>. Another study found that the contribution of obesity and craniofacial features to OSA differed between Caucasians and Chinese patients with similar OSA severity- the Caucasians were more obese while the Chinese exhibited more craniofacial restriction<sup>17;18</sup>. However, whether non-anatomical factors contribute to the ethnic differences in OSA phenotype has not previously been studied.

In this issue of *Respirology* Lee et al. provide evidence that non-anatomical factors contributing to the pathogenesis of OSA differ significantly between White Caucasians and Chinese with moderate to severe OSA (based on polysomnography). This study shows that a low respiratory ArTH is significantly less common in Chinese patients with moderate to severe OSA compared to White Caucasians and that the ArTH was less negative in Caucasians compared to Chinese with moderate to severe OSA<sup>19</sup>. Low respiratory ArTH in this study was defined as having at least two of the following criteria: apnoea hypopnea index < 30/hour, nadir oxygen saturation > 82.5%, and percentage of hypopneas out of the total respiratory events > 58.3%<sup>19</sup>. This scoring system was validated to predict a low ArTH (defined as overall ArTH less negative than -15 cmH<sub>2</sub>O on epiglottic pressure catheter) by Bradley et al.; a score of 2 or more had correctly predicted a low ArTH in 84.1% of cases with a sensitivity, a specificity, a positive predictive value and a negative predictive value all over 80%<sup>11</sup>. In addition, and after considering craniofacial anatomy, the proportion of White Caucasians with moderate to severe OSA who had low respiratory ArTH was lower in those who had severe craniofacial anatomical compromise; While in the Chinese patients with moderate to severe OSA there was no such differences in the prevalence of low respiratory ArTH between those with and without craniofacial anatomical compromise<sup>19</sup>. In a sub-group analysis in which the White Caucasian and Chinese patients with moderate-severe OSA were matched for BMI; OSA was more

severe in the Chinese patients and the ArTH was less negative in the White Caucasians. However, matching for BMI does not necessarily reflect matching for adiposity as Asians are known to have higher visceral adiposity compared to White Caucasians with similar BMI<sup>20</sup>. This might explain the higher prevalence of OSA in the Chinese patients compared to the White Caucasians with similar BMI in this subgroup analysis. It must be noted that the criteria used for assessing low ArTH in this study were validated previously in White Caucasian populations and not in Chinese population<sup>11</sup>.

This study suggests that OSA in Chinese patients is more driven by anatomical factors while in White Caucasians non-anatomical factors play a more important role. This is relevant when considering individualising screening and treatment strategies for patients with OSA. For example, treatments aimed at improving low respiratory ArTH may only be beneficial in patients who have low respiratory ArTH as part of the pathogenesis of their OSA, while in other patients such treatment may have no impact or even a detrimental effect on AHI<sup>21;22</sup>.

In conclusion, the current study by Lee et al suggests that there are ethnic differences in the pathogenesis of OSA between Chinese and White Caucasians with moderate to severe OSA. The authors demonstrate that the contributions of obesity, craniofacial anatomy and respiratory ArTH to OSA pathogenesis differs between White Caucasians and Chinese patients with moderate to severe OSA. Further research exploring ethnic differences in OSA pathogenesis is needed in order to individualise treatment and screening strategies.

## **Acknowledgment**

Abd Tahrani is a clinician scientist supported by the National Institute for Health Research in the UK. The views expressed in this publication are those of the author(s) and not necessarily those of the National Health Service, the National Institute for Health Research or the Department of Health.

## Reference List

- (1) Young T, Peppard PE, Gottlieb DJ. Epidemiology of Obstructive Sleep Apnea: A Population Health Perspective. *Am J Respir Crit Care Med* 2002; 165(9):1217-1239.
- (2) McNicholas WT. Diagnosis of Obstructive Sleep Apnea in Adults. *Proceedings of the American Thoracic Society* 2008; 5(2):154-160.
- (3) Tahrani AA, Ali A, Stevens MJ. Obstructive sleep apnoea and diabetes: an update. *Current Opinion in Pulmonary Medicine* 2013; 19(6).
- (4) Newman AB, Foster G, Givelber R, Nieto FJ, Redline S, Young T. Progression and Regression of Sleep-Disordered Breathing With Changes in Weight: The Sleep Heart Health Study. *Arch Intern Med* 2005; 165(20):2408-2413.
- (5) Greenburg DL, Lettieri CJ, Eliasson AH. Effects of Surgical Weight Loss on Measures of Obstructive Sleep Apnea: A Meta-Analysis. *Am J Med* 2009; 122(6):535-542.
- (6) Shigeta Y, Ogawa T, Tomoko I, Clark GT, Enciso R. Soft palate length and upper airway relationship in OSA and non-OSA subjects. *Sleep Breath* 2010; 14(4):353-358.
- (7) Lee RWW, Sutherland K, Chan ASL, Zeng B, Grunstein RR, Darendeliler MA et al. Relationship Between Surface Facial Dimensions and Upper Airway Structures in Obstructive Sleep Apnea. *Sleep* 2010; 33(9):1249-1254.
- (8) Sutherland K, Lee RWW, Petocz P, Chan TO, Ng S, Hui DS et al. Craniofacial phenotyping for prediction of obstructive sleep apnoea in a Chinese population. *Respirology* 2016; 21(6):1118-1125.
- (9) Lam B, Ip MSM, Tench E, Ryan CF. Craniofacial profile in Asian and white subjects with obstructive sleep apnoea. *Thorax* 2005; 60(6):504.
- (10) Eckert DJ, White DP, Jordan AS, Malhotra A, Wellman A. Defining Phenotypic Causes of Obstructive Sleep Apnea. Identification of Novel Therapeutic Targets. *Am J Respir Crit Care Med* 2013; 188(8):996-1004.
- (11) Edwards BA, Eckert DJ, McSharry DG, Sands SA, Desai A, Kehlmann G et al. Clinical Predictors of the Respiratory Arousal Threshold in Patients with Obstructive Sleep Apnea. *Am J Respir Crit Care Med* 2014; 190(11):1293-1300.
- (12) Tahrani AA. Diabetes and sleep apnea. *International Textbook of Diabetes Mellitus*. John Wiley & Sons, Ltd; 2015. 316-336.
- (13) Chen X, Wang R, Zee P, Lutsey PL, Javaheri S, Alcaíntara C et al. Racial/Ethnic Differences in Sleep Disturbances: The Multi-Ethnic Study of Atherosclerosis (MESA). *Sleep* 2015; 38(6):877-888.
- (14) Villaneuva AT, Buchanan PR, Yee BJ, Grunstein RR. Ethnicity and obstructive sleep apnoea. *Sleep Medicine Reviews* 2005; 9(6):419-436.

- (15) Punjabi NM. The Epidemiology of Adult Obstructive Sleep Apnea. *Proc Am Thorac Soc* 2008; 5(2):136-143.
- (16) Amin A, Ali A, Altaf QA, Piya MK, Barnett AH, Raymond NT et al. Prevalence and Associations of Obstructive Sleep Apnea in South Asians and White Europeans with Type 2 Diabetes: A Cross-Sectional Study. *Journal of clinical sleep medicine: JCSM: official publication of the American Academy of Sleep Medicine*. In press 2017.
- (17) Lee RWW, Vasudavan S, Hui DS, Prvan T, Petocz P, Darendeliler MA et al. Differences in Craniofacial Structures and Obesity in Caucasian and Chinese Patients with Obstructive Sleep Apnea. *Sleep* 2010; 33(8):1075-1080.
- (18) Li KK, Kushida C, Powell NB, Riley RW, Guilleminault C. Obstructive Sleep Apnea Syndrome: A Comparison Between Far-East Asian and White Men. *The Laryngoscope* 2000; 110(10):1689-1693.
- (19) Lee RWW, Sutherland K, Sands SA, Edwards BA, Chan TO, Ng S et al. Differences in Respiratory Arousal Threshold in Caucasians and Chinese Patients with Obstructive Sleep Apnoea. *Respirology*. In press 2017.
- (20) Gray LJ, Yates T, Davies MJ, Brady E, Webb DR, Sattar N et al. Defining Obesity Cut-Off Points for Migrant South Asians. *PLoS ONE* 2011; 6(10):e26464.
- (21) Carter SG, Berger MS, Carberry JC, Bilston LE, Butler JE, Tong BK et al. Zopiclone Increases the Arousal Threshold without Impairing Genioglossus Activity in Obstructive Sleep Apnea. *Sleep* 2016; 39(4):757-766.
- (22) Eckert DJ, Owens RL, KEHLMANN GB, Wellman A, RAHANGDALE S, YIM-YEH S et al. Eszopiclone increases the respiratory arousal threshold and lowers the apnoea/hypopnoea index in obstructive sleep apnoea patients with a low arousal threshold. *Clin Sci (Lond)* 2011; 120(12):505-514.